



ORAL PRESENTATION

Open Access

MRI detects coronary vessel wall thickening with age in healthy subjects

Andrew D Scott^{1*}, Jennifer Keegan², Raad H Mohiaddin², David N Firmin¹

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Objective

To investigate the effects of ageing on the coronary vessel wall of healthy subjects using MR.

Background

Autopsy studies [1,2] and x-ray CT in older subjects with suspected disease [3] have demonstrated increasing coronary vessel thickness with age. There is, however, a need for a radiation-free non-invasive technique for use in longitudinal studies of coronary vessel wall morphology. Recently 3D MR coronary vessel wall imaging with retrospective beat-to-beat respiratory-motion-correction (B2B-RMC)[4], which uses 3D low resolution data acquired immediately before the main imaging data to retrospectively correct respiratory motion, has demonstrated great promise [5]. We propose that 3D high resolution MR with B2B-RMC can

demonstrate coronary vessel wall thickening with age in healthy subjects.

Methods

21 healthy subjects with no history of cardiovascular disease (mean age 39 ± 13 , range 22-62, 11 female) were recruited. Studies were performed on a 1.5T Siemens Avanto scanner. Cross-sectional vessel wall imaging was performed in the proximal right coronary artery (<40mm from origin) using a 3D spiral acquisition with B2B-RMC (0.7x0.7x3mm resolution, 8 slices reconstructed to 16x1.5mm, duration 600 cardiac cycles assuming 100% respiratory efficiency) and alternate R-wave cardiac gating. Data were acquired in the coronary rest period, as determined from a cine acquisition in the imaging plane. Circular regions of interest were drawn around the inner and outer coronary vessel wall on one




Figure 1 Example images obtained from three male healthy subjects aged 24 (wall thickness 1.02mm, W/OW 0.69) (a), aged 40 (wall thickness 1.38mm, W/OW 0.73) (b) and aged 59 (wall thickness 1.53mm, W/OW 0.70) (c).

¹Imperial College London, London, UK

Full list of author information is available at the end of the article




Figure 2 The effects of ageing on the coronary vessel wall as assessed by high resolution MR imaging. Vessel wall thickness (a) and W/OW (b) demonstrate a significant positive correlation with age.

central slice from each 3D acquisition. Average vessel wall thickness and wall/outer wall (lumen + vessel wall) area (W/OW)[6] were calculated.

Results

Example images are shown in figure 1. In three subjects (14%) the images were rejected due to poor image quality caused by cardiac or respiratory motion. In the remaining 18 subjects, mean vessel wall thickness was 1.14 ± 0.22 mm and mean W/OW was 0.727 ± 0.085 . Vessel wall thickness and W/OW increase by 0.088 mm ($R=0.53, p=0.024$, figure 2.a) and 0.031 ($R=0.48, p=0.048$, figure 2.b) per decade respectively.

Discussion

For the first time using MR, we have demonstrated significantly increasing vessel wall thickness with age in a small cohort of healthy subjects at 0.088 mm per decade. The proportion of vessel that is vessel wall (W/OW) also increases with age at 0.031 per decade ($R=0.48$). The strength of this correlation is similar to that obtained in larger studies of carotid wall thickening in healthy subjects ($R=0.50$ males and $R=0.46$ females [7]).

Author details

¹Imperial College London, London, UK. ²The Royal Brompton and Harefield NHS Foundation Trust, London, UK.

Published: 2 February 2011

References

1. Velican: *Atherosclerosis* 1980.
2. Ozolanta: *Med Eng Phys* 1998.
3. Matsunaga: *Circulation J* 2009.
4. Keegan: *JMRI* 2007.
5. Scott: *JMRI*.
6. Saam: *JCMR* 2005.
7. Keenan: *Atherosclerosis* 2009.

doi:10.1186/1532-429X-13-S1-O47

Cite this article as: Scott et al.: MRI detects coronary vessel wall thickening with age in healthy subjects. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):O47.