

ORAL PRESENTATION

Open Access

Impaired myocardial perfusion on CMR is associated with increased numbers of classical monocytes in STEMI patients treated by primary PCI

Lourens FHJ Robbers^{1*}, Anja M van der Laan², Robin Nijveldt¹, Aernout M Beek¹, Ronak Delewi¹, Alexander Hirsch², C Ellen van der Schoot³, Bart J Biemond², Felix Zijlstra⁴, Jan J Piek², Albert C van Rossum¹

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Introduction

Despite optimal revascularization strategies, STEMI patients often show impaired myocardial tissue reperfusion due to microvascular obstruction. Monocytes, especially the classical monocytes have been suggested to contribute to microvascular obstruction through a complex inflammatory response.

Purpose

To investigate the relation between the circulating number of classical monocytes and severity of myocardial perfusion impairment in patients with ST-elevated myocardial infarction (STEMI) treated with primary PCI.

Methods


Fifty-four patients underwent CMR first-pass perfusion imaging, late gadolinium enhancement (LGE) and functional imaging between 3-7 days after primary PCI.

Myocardial perfusion in the infarct core was evaluated semi-quantitatively from signal intensity versus time curves by calculation of the maximum upslope relative to a remote, normal segment. The number of classical ($CD14^{++}CD62L^{+}$) (CM) and nonclassical ($CD14^{+}CD62L^{-}$) monocytes (NCM) was measured by flow cytometry from a venous blood sample. Patients were divided in tertiles according to the monocyte count in the sample.

Results

Myocardial perfusion in the infarct core was significantly lower in patients with high numbers of CM (upper

tertile) compared to low numbers (lower tertile) of CM (mean relative upslope in upper tertile 7.5 versus mid tertile 5.8 versus lower tertile 5.2, $p=0.014$). Also, patients with high numbers of CM had a larger infarct size (9.17 vs. 6.51 vs. 3.89 g/m^2 , $p<0.001$), more segments with microvascular obstruction at LGE (15.9 vs. 8.2 vs. 2.2% of segments, $p=0.001$), and greater impairment of ejection fraction (37.0 vs. 42.0 vs. 46.5%, $p=0.001$) than patients with low numbers of CM. No such relations were found for NCM. Figure 1.



¹VU University Medical Center, Amsterdam, Netherlands
Full list of author information is available at the end of the article

Conclusions

In reperfused STEMI patients, high numbers of classical monocytes are associated with impaired myocardial perfusion, increased myocardial injury and greater functional loss. This suggests a causal relationship between circulating monocytes after AMI and degree of myocardial damage in reperfused infarction.

Author details

¹VU University Medical Center, Amsterdam, Netherlands. ²Academic Medical Center, Amsterdam, Netherlands. ³Sanquin Research and Landsteiner Laboratory, Amsterdam, Netherlands. ⁴University Medical Center Groningen, Groningen, Netherlands.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-O93

Cite this article as: Robbers *et al.*: Impaired myocardial perfusion on CMR is associated with increased numbers of classical monocytes in STEMI patients treated by primary PCI. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):O93.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

