

POSTER PRESENTATION

Open Access

Effectiveness of cardioprotective medication in women with suspected ischemic heart disease syndrome: the NHLBI-sponsored women's ischemia syndrome evaluation (WISE) study

Mark Doyle^{1*}, Gerald Pohost², Leslie Shaw³, Sheryl Kelsey⁴, Diane Vido¹, Delia Johnson⁴, William J Rogers⁵, William J Rogers⁵, Geetha Rayarao¹, Barry Sharaf², Carl J Pepine⁶, C Noel Bairey Merz³, Robert WW Biederman¹

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Introduction

Knowledge of the effectiveness of cardioprotective medication in women with suspected ischemia is hampered by limited clinical trial data and heterogeneous risk. We assessed the effectiveness of cardioprotective medical treatment in women while adjusting for disease severity determined by cardiovascular magnetic resonance imaging (CMRI).

Hypothesis

Effectiveness of cardioprotective medication will be evident in women at elevated risk determined by CMRI.

Methods

Women (n=113), mean age 58±12 years, undergoing coronary angiography for symptoms suggestive of myocardial ischemia additionally underwent myocardial perfusion and cardiac function CMRI evaluation. Previously we identified four ischemic heart disease components associated with adverse cardiac events: 1) coronary artery stenosis > 50%, 2) low global myocardial perfusion, 3) high cardiac energy utilization, and 4) myocardial wall thickness > 10 mm. The latter three disease components were measured using CMRI. Women were stratified into 5 groups based on number of ischemic heart disease components (0-4). Self-reported medication use included angiotensin converting enzyme inhibitors, beta blockers (BB), calcium channel antagonists (CCA) and nitrates. During follow-up (32±17 months)

time to first adverse event (death, myocardial infarction, hospitalization for either congestive heart failure or for worsening anginal symptoms) was analyzed. Cox proportional hazard regression was performed to examine the cardioprotective effectiveness of medications when accounting for ischemic heart disease severity.

Results

Medication usage did not differ across ischemic heart disease strata. Ischemic heart disease stratification predicted adverse cardiac events (hazard ratio 2.1, 95% confidence interval 1.5-2.9, p<0.001) and the medication combination of a BB or CCA and no nitrate were independently associated with reduced risk of events (hazard ratio 0.35, 95% confidence interval 0.14-0.86, p<0.05). The model remained significant after adjustment for age and Framingham risk score.

Conclusions

Using a combined model of ischemic heart disease severity and medication use, the medication combination of beta-blocker or calcium antagonist and no nitrate were associated with a decreased risk of adverse cardiac events in women. Further investigation of these results should be done with a prospective randomized clinical trial.

Author details

¹Allegeny General Hospital, Pittsburgh, PA, USA. ²Brown University, Providence, RI, USA. ³Cedars-Sinai Medical Center, Los Angeles, CA, USA. ⁴University of Pittsburgh, Pittsburgh, PA, USA. ⁵University of Alabama, Birmingham, AL, USA. ⁶University of Florida, Gainesville, FL, USA.

¹Allegeny General Hospital, Pittsburgh, PA, USA
Full list of author information is available at the end of the article

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P169

Cite this article as: Doyle *et al.*: Effectiveness of cardioprotective medication in women with suspected ischemic heart disease syndrome: the NHLBI-sponsored women's ischemia syndrome evaluation (WISE) study. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P169.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

