

POSTER PRESENTATION

Open Access

# Regional longitudinal bi-ventricular function in pulmonary hypertension: single heart-beat assessment of strain by fast-senc imaging

Monda L Shehata<sup>1\*</sup>, Tamer A Basha<sup>1</sup>, Ahmed A Harouni<sup>1</sup>, Jan Skrok<sup>1</sup>, Sukhminder Singh<sup>1</sup>, Stephen Mathai<sup>1</sup>, Joao AC Lima<sup>1</sup>, Nael F Osman<sup>1</sup>, David A Bluemke<sup>2</sup>, Paul M Hassoun<sup>1</sup>, Jens Vogel-Claussen<sup>1</sup>


From 2011 SCMR/Euro CMR Joint Scientific Sessions  
Nice, France. 3-6 February 2011

## Purpose

Right ventricular (RV) function is the most important determinant of survival in patients with pulmonary hypertension (PH), thus monitoring of RV function is critical. However, RV regional function assessment is challenging using current MR tagging techniques. Thus, the purpose was to evaluate regional longitudinal ventricular deformation ( $E_{LL}$ ) acquired by free breathing single heart-beat fast strain encoded imaging (fast-SENC) in relation to global ventricular dysfunction markers and pulmonary hemodynamics in PH patients.

## Materials and methods

48 subjects [35 PH patients (mean pulmonary arterial pressure mPAP =  $40.2 \pm 11.8$  mmHg) and 13 age and gender matched controls] were examined using short axis fast-SENC MRI and cine imaging. All patients underwent right heart catheterization (RHC). Segmental (15 RV and 16 LV segments), slice (basal, mid and apical), as well as mean (average of segments) peak systolic  $E_{LL}$  were quantified for both ventricles and correlated with global function and RHC indices. Patients were stratified into 3 groups based on RV ejection fraction (RVEF).




**Figure 1** (A) Basal short axis fast-SENC peak systolic image in a 53 year old healthy female subject demonstrating normal longitudinal shortening ( $E_{LL}$ ) (red color). (B) Basal short axis fast-SENC peak systolic image in a 47 year old female PH patient (mPAP = 51 mmHg). Note regionally reduced RV  $E_{LL}$  and LV antero-septal  $E_{LL}$  (white color). Probe points correspond to regions of interest where strain was measured. A mesh was used to measure LV  $E_{LL}$  at six basal, six mid and four apical LV segments respectively.

<sup>1</sup>Johns Hopkins University, Baltimore, MD, USA  
Full list of author information is available at the end of the article

**Table 1 Right ventricular longitudinal (ELL) strain in PH patients and controls**

ELL (%)	PH 25% - 75% (N=35)	Control 25% - 75% (N =13)	P value
Mean Ventricular Strain	-19.3 (-20.5 - -15.3)	-21.3 (-21.9 - -20.2)	0.001
Slices			
Basal	-19.1 (-21.6 - -14.9)	-21.7 (-22.0 - -20.2)	0.008
Mid	-19.3 (-20.3 - -12.6)	-21.8 (-22.7 - -19.1)	0.002
Apical	-18.1 (-21.8 - -15.7)	-20.9 (-22.3 - -19.8)	0.02
Segments			
Basal Anterior Septal Insertion	-14.1 (-18.8 - -12.5)	-20.2 (-23.1 - -17.1)	<0.001
Basal Anterior	-21.2 (-24.1 - -15.1)	-24.0 (-25.1 - -21.8)	0.02
Basal Lateral	-21.0 (-25.1 - -17.9)	-23.8 (-26.2 - -22.5)	0.03
Basal Inferior	-18.3 (-21.6 - -15.1)	-20.2 (-23.6 - -18.6)	0.09
Basal Inferior Septal Insertion	-19.9 (-21.6 - -14.1)	-19.1 (-21.7 - -16.3)	0.88
Mid Anterior Septal Insertion	-15.1 (-18.1 - -9.6)	-20.8 (-22.7 - -17.7)	<0.001
Mid Anterior	-16.3 (-19.1 - -12.5)	-22.1 (-25.1 - -21.3)	<0.001
Mid Lateral	-22.5 (-23.6 - -15.0)	-24.9 (-25.9 - -22.9)	0.005
Mid Inferior	-20.1 (-23.3 - -14.1)	-20.1 (-24.6 - -18.7)	0.16
Mid Inferior Septal Insetion	-16.9 (-21.2 - -11.1)	-18.7 (-20.1 - -16.1)	0.49
Apical Anterior Septal Insertion	-16.6 (-20.7 - -12.3)	-20.4 (-22.3 - -19.0)	0.02
Apical Anterior	-18.4 (-21.2 - -14.9)	-21.9 (-23.8 - -18.8)	0.01
Apical Lateral	-20.4 (-24.4 - -15.4)	-24.1 (-25.7 - -22.1)	0.0009
Apical Inferior	-18.3 (-24.1 - -13.1)	-22.2 (-24.6 - -19.7)	0.08
Apical Inferior Septal Insertion	-18.1 (-21.8 - -12.8)	-17.3 (-19.5 - -15.6)	0.84



**Figure 2**

## Results

PH patients demonstrated reduced  $E_{LL}$  at all RV levels compared to controls (Figure 1). On regional analysis, reduced  $E_{LL}$  was noted in all segments ( $p < 0.05$ ) except the inferior and inferior septal insertion at all levels (Table 1). Reduced mean RV  $E_{LL}$  correlated with elevated mPAP ( $r = 0.6$ ,  $p < 0.001$ ), pulmonary vascular resistance index (PVRI,  $r = 0.8$ ,  $p < 0.001$ ) as well as reduced RV systolic function parameters ( $p < 0.05$  for all) (Figure 2). In the LV, reduced  $E_{LL}$  was mainly noted at the basal anterior (-16.7 vs. -20.5,  $p = 0.03$ ) and antero-septal regions (-12.8 vs. -18.2,  $p < 0.01$ ). Reduced LV antero-septal  $E_{LL}$  correlated with increased mPAP ( $r = 0.5$ ,  $p < 0.01$ ), increased septal eccentricity index ( $r = 0.5$ ,  $p < 0.01$ ) and reduced RV systolic function ( $p < 0.05$  for all). On multiple linear regression including mPAP, RV end-diastolic volume index and RV mass index as covariates, mPAP was an independent predictor of reduced mean RV  $E_{LL}$  ( $\beta = 0.19$ ,  $p < 0.01$ ). In turn, reduced mean RV  $E_{LL}$  was the main predictor of reduced RVEF in PH patients ( $\beta = -1.2$ ,  $p = 0.03$ ) in a model including mPAP, mean RV  $E_{LL}$  and RV mass index. In PH patients with maintained global RV function, regional  $E_{LL}$  was reduced at the basal and mid anterior septal insertions as well as mid anterior RV segments ( $p < 0.05$  for all).

## Conclusion

In PH patients, reduced RV  $E_{LL}$  measured by fast-SENC is associated with increased afterload and correlates with biventricular global dysfunction. RV strain analysis using fast-SENC can detect subclinical regional dysfunction in absence of global RV functional compromise.

### Author details

<sup>1</sup>Johns Hopkins University, Baltimore, MD, USA. <sup>2</sup>National Institutes of Health, Bethesda, MD, USA.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P279

**Cite this article as:** Shehata et al.: Regional longitudinal bi-ventricular function in pulmonary hypertension: single heart-beat assessment of strain by fast-senc imaging. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P279.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

