

POSTER PRESENTATION

Open Access

# Effect of uphill running on myocardium $T_2$ in *mdx* mice

Sean C Forbes<sup>1\*</sup>, Ravneet S Vohra<sup>1</sup>, Fan Ye<sup>1</sup>, Krista Vandendorpe<sup>1</sup>, Glenn A Walter<sup>2</sup>

From 15th Annual SCMR Scientific Sessions  
Orlando, FL, USA. 2-5 February 2012

## Background

Cardiac dysfunction is a major cause of death in Duchenne muscular dystrophy. In *mdx* mice, the lack of functional dystrophin localized to the cell membrane leads to increased susceptibility to muscle damage and enhanced muscle degeneration. In this study we examined the effect of an uphill running protocol (Michele et al. *Circ Res.* 105(10):984-93, 2009) on myocardium transverse relaxation time ( $T_2$ ) in young adult *mdx* mice (16 weeks).

## Methods

A 4.7T Oxford Magnet with an Agilent/Varian operating system was used to acquire gated  $T_2$ -weighted single spin-echo images of the left ventricle in the short axis view (TR 750 ms; TE 14-16 ms and TE 30-32 ms; field of view, 25X25 mm<sup>2</sup>; slice thickness, 1.0 mm; acquisition matrix size, 256 X 128; averages, 8). Images were acquired in C57Bl10 (n=5, male) and *mdx* (n=5, male) mice using a custom built quadrature volume coil (3.3 cm inner diameter). Mice performed uphill treadmill running at a speed of 6-13m/min with a 10 degree incline for up to one hour. MR data were acquired prior to exercise and after 16-24 hrs of exercise. Short axis slices from the mid-papillary region were selected to calculate mean  $T_2$ . Mean  $T_2$  of the myocardium was calculated using the average signal intensity at each TE by manually tracing the myocardium.


## Results

Each control mouse completed one hour of uphill running (800 meters), while there was considerable variability in the amount of time and distance run by the *mdx* mice (33±21(SD) min; 329±227 meters), with only one *mdx* mouse completing one hour. In wild-type mice,

there was no effect ( $p>0.05$ ) of this uphill running protocol on myocardium  $T_2$  after exercise (Fig. 1). In *mdx* mice, an increase in myocardium  $T_2$  was observed in each mouse (range: 5 to 32%; Fig. 1). There did not appear to be a direct relationship between running time and  $T_2$  increase ( $r=.14$ ,  $p>0.05$ ).

## Conclusions

The increase in myocardium  $T_2$  following exercise in *mdx* mice is consistent with dystrophic muscle having an increased susceptibility to muscle damage. Therefore, this *in vivo* exercise protocol monitored using cardiac MRI may be valuable in future studies to test the efficacy of potential therapeutic treatments in dystrophic murine models. Furthermore, this study supports the notion that  $T_2$  may be valuable for evaluating myocardium involvement in muscular dystrophy.



**Figure 1** Myocardium  $T_2$  values before and after uphill running in wild-type and *mdx* mice. \*Denotes significantly different than pre-exercise.

<sup>1</sup>Department of Physical Therapy, University of Florida, Gainesville, FL, USA  
Full list of author information is available at the end of the article

## Funding

Supported by American Heart Association Post-doctoral Fellowship and Wellstone Muscular Dystrophy Center (1U54RO52646-01A1).

## Author details

<sup>1</sup>Department of Physical Therapy, University of Florida, Gainesville, FL, USA.

<sup>2</sup>Physiology and Functional Genomics, University of Florida, Gainesville, FL, USA.

Published: 1 February 2012

doi:10.1186/1532-429X-14-S1-P59

**Cite this article as:** Forbes et al.: Effect of uphill running on myocardium T<sub>2</sub> in *mdx* mice. *Journal of Cardiovascular Magnetic Resonance* 2012 **14** (Suppl 1):P59.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

