

ORAL PRESENTATION

Open Access

Diagnostic accuracy of the core components of a multi-parametric cardiovascular magnetic resonance imaging protocol: a CE-MARC sub-study

John P Greenwood^{1*}, David P Ripley¹, Manish Motwani¹, Julia Brown², Jane Nixon², Colin C Everett², Neil Maredia¹, Petra Bijsterveld¹, Sven Plein¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

The CE-MARC study was the largest prospective evaluation of the diagnostic accuracy of cardiovascular magnetic resonance (CMR) in coronary artery disease (CAD), demonstrating its superiority over single-photon emission computed tomography [1]. The trial adopted a multi-parametric protocol assessing ventricular function, myocardial perfusion, viability (with late gadolinium enhancement (LGE)) and coronary artery anatomy. There have been a number of previous studies analysing the diagnostic accuracy of different components of the CMR examination with contrasting results. We assessed the diagnostic accuracy of the individual components and selected combinations of the multi-parametric CMR examination from the CE-MARC study.

Methods

All patients from the CE-MARC population were studied. Visual CMR analyses were from the original, blinded read. Pre-specified sub-analysis of the four individual core components of the CMR protocol was performed in isolation, as a paired component (perfusion and LGE) and as a triplet (perfusion, LGE and ventricular function) and compared to the full multi-parametric protocol.

Results

Both CMR and X-ray angiography were available in 676 patients. The sensitivity of the combined CMR protocol

was 86.5%, specificity 83.4%, PPV 77.2% and NPV 90.5%. The diagnostic accuracy of the individual components and paired and triplet combinations compared to the full multi-parametric protocol are presented in Table 1 and Figure 1. The maximum sensitivity for the detection of significant CAD by CMR was achieved when all four components were used. No individual component; paired perfusion with LGE; or perfusion with LGE and function significantly outperformed the multi-parametric protocol in terms of sensitivity. However in terms of specificity, the individual components of perfusion, ventricular function and late gadolinium enhancement (LGE) all performed significantly better than the multi-parametric protocol ($P < 0.0001$). In addition, combining LGE with perfusion or with perfusion and ventricular function significantly improved the test specificity compared to the multi-parametric protocol ($P < 0.0001$ for each). In terms of PPV and NPV, the multi-parametric protocol performed significantly better than all individual components, paired or triplet combination.

Conclusions

A combined multi-parametric CMR protocol has higher sensitivity, PPV and NPV than the individual components however the specificity of the individual components of perfusion, ventricular function and late gadolinium enhancement (LGE) all performed significantly better than the multi-parametric protocol.

Funding


This study was funded by the British Heart Foundation (RG/05/004). J.P.G and S.P. received research grant

¹Multidisciplinary Cardiovascular Research Centre (MCRC) & Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, Leeds, UK
Full list of author information is available at the end of the article

Table 1 Diagnostic accuracy of a multi-parametric CMR exam and its core components compared to the reference test X-ray angiography.

	Sensitivity (95%CI)	Specificity (95%CI)	PPV (95%CI)	NPV (95%CI)
Overall multi-parametric CMR study (all components) (n = 676)	86.5 (81.8, 90.1)	83.4 (79.5, 86.7)	77.2 (72.1, 81.6)	90.5 (87.1, 93.0)
Single CMR components				
LGE (n = 674)	40.8 (35.0, 46.8)	95.8 (93.4, 97.4)	86.4 (79.3, 91.3)	71.4 (67.5, 75.0)
Perfusion (n = 661)	76.9 (71.4, 81.6)	91.8 (88.7, 94.1)	85.8 (80.8, 89.7)	86.0 (82.4, 89.0)
Ventricular function (n = 676)	47.4 (41.4, 53.4)	93.7 (90.9, 95.6)	82.9 (76.1, 88.1)	73.3 (69.3, 76.9)
MRA (n = 597) **	71.2 (65.1, 76.7)	89.8 (86.3, 92.5)	81.8 (75.9, 86.5)	83.0 (79.0, 86.4)
Combinations				
Perfusion/LGE (n = 676)	78.6 (73.3, 83.1)	89.3 (85.9, 91.9)	82.6 (77.5, 86.8)	86.5 (82.9, 89.5)
Perfusion/LGE/ventricular function (n = 676)	81.6 (76.5, 85.8)	85.9 (82.1, 88.9)	78.9 (73.7, 83.3)	87.8 (84.2, 90.6)

CMR - cardiovascular magnetic resonance; LGE - late gadolinium enhancement; MRA - magnetic resonance angiography. ** Only includes excellent or adequate quality MRA



from Philips Healthcare. S.P. is funded by a British Heart Foundation fellowship (FS/10/62/28409).

Authors' details

¹Multidisciplinary Cardiovascular Research Centre (MCRC) & Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, Leeds, UK. ²Clinical Trials Research Unit, University of Leeds, Leeds, UK.

Published: 16 January 2014

Reference

1. Greenwood JP, Maredia N, Younger JF, Brown JM, Nixon J, Everett CC, *et al.*: Cardiovascular magnetic resonance and single-photon emission computed tomography for diagnosis of coronary heart disease (CE-MARC): a prospective trial. *Lancet* 2012, **379**(9814):453-60.

doi:10.1186/1532-429X-16-S1-O19

Cite this article as: Greenwood *et al.*: Diagnostic accuracy of the core components of a multi-parametric cardiovascular magnetic resonance imaging protocol: a CE-MARC sub-study. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):O19.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

