

POSTER PRESENTATION



A new method for vessel segmentation based on a priori input from medical expertise in cine phase-contrast Magnetic Resonance Imaging

Sebastian L Bidhult^{1*}, Marcus Carlsson¹, Katarina Steding-Ehrenborg¹, Håkan Arheden¹, Einar Heiberg^{1,2}

From 17th Annual SCMR Scientific Sessions New Orleans, LA, USA. 16-19 January 2014


Background

Phase contrast magnetic resonance imaging (PC-MRI) is the current gold standard for blood flow quantification. Using this technique it is possible to quantify cardiac output, stroke volume, shunts and valve insufficiencies. The accuracy of flow quantification is highly dependent on the manual delineation of the vessels of interest. Therefore, the purpose of this study is to develop an automatic method for vessel segmentation in PC-MRI sequences.

Methods

A total of 211 subjects from a previous study [1] were included (160 heart failure patients with ejection fraction below 40% and 51 healthy volunteers). For all subjects

manual delineations of aorta ascendens were available in all timeframes. The subjects were randomly divided into a training (n = 40) and a test set (n = 171). The manual delineations in the training set were parameterized and principle component analysis (PCA) was performed to extract a priori information regarding vessel shape change and motion during the cardiac cycle. The proposed algorithm is an active contour model which tracks the edges of the aortic wall, initialized by a manual delineation in one timeframe. Every deformation is followed by a projection onto a subset of linearly independent components from the PCA result of the a priori dataset. This procedure is repeated 8 times. Validation was performed by comparing automated delineations with manual delineations in the test set consisting of the 171 remaining patients.



¹Department of Clinical Physiology, Lund University, Lund, Skane, Sweden Full list of author information is available at the end of the article



© 2014 Bidhult et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http:// creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

All automatic segmentations were initiated with the manual delineation from the user in one timeframe. Stroke volume was used as performance measure. The inter observer variability for manual delineations from two observers was calculated in the previous study [1].

Results

The stroke volume from the automated method agreed well with manual delineations (Figure 1), with a correlation coefficient of r2 = 0.98. Bias between automated and manual delineation was -0.69 ± 2.55 ml, or $-0.7 \pm 3.5\%$ of the stroke volume. This should be compared to an inter observer variability of $3 \pm 4\%$ [1]. The low bias of the proposed method compared to the inter observer variability of the previous study [1] is likely attributed to the fact that the initialization of the automatic method was done by the observer to whom the comparison was made.

Conclusions

In conclusion, the proposed method for vessel segmentation results in bias and variability comparable to manual delineations.

Funding

Swedish Research Council, Swedish Heart-Lung Foundation, Medical Faculty at Lund University.

Authors' details

¹Department of Clinical Physiology, Lund University, Lund, Skane, Sweden. ²Centre for Mathematical Sciences, Lund University, Lund, Skane, Sweden.

Published: 16 January 2014

Reference

1. Carlsson M, et al: JCMR 2012, 14(51).

doi:10.1186/1532-429X-16-S1-P355 Cite this article as: Bidhult *et al*: A new method for vessel segmentation based on a priori input from medical expertise in cine phase-contrast Magnetic Resonance Imaging. *Journal of Cardiovascular Magnetic Resonance* 2014 16(Suppl 1):P355.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

BioMed Central