

POSTER PRESENTATION



Arterial spin labeling CMR perfusion imaging is capable of continuously monitoring myocardial blood flow during stress

Hung P Do^{1,2*}, Ahsan Javed², Terrence R Jao³, Hee-won Kim⁴, Andrew J Yoon⁵, Krishna S Nayak²

From 18th Annual SCMR Scientific Sessions Nice, France. 4-7 February 2015

Background

Myocardial arterial spin labeling (ASL) is a non-invasive non-contrast CMR perfusion imaging technique [Zun et al., '09] that is compatible with adenosine stress testing [Zun et al., '11]. Here, we demonstrate its ability to continuously monitor MBF during stress, which has a variety of potential applications including interventional CMR and the study of endothelial function. We used mild sustained isometric handgrip stress.

Methods

Imaging methods

Three healthy subjects (25-30, 3 males, non-smoker, no family history of CAD) were scanned on a GE 3T scanner. A previously described myocardial ASL sequence [Zun et al., '09; Zun et al., '11; Do et al., '14] was run continuously for roughly 20 minutes, in a free-breathing mode with respiratory triggering, on a single mid short-axis slice. The imaging protocol was divided into three stages: 6 min rest, 6 min stress, and 6-8min recovery. Imaging parameters were TE/TR=1.5/3.2ms, FOV = 240-320mm, matrix size = 96x96, slice thickness = 10mm, and an acceleration factor of 2. Stress was induced using a sustained isometric handgrip exercise at 30% of each subject's maximum voluntary contraction [Brenkenhoff et al., '13].

Data analysis

Motion correction [Avants et al., '08] was used to generate myocardial masks of all acquired control and tagged image pairs from a manually segmented reference pair [Javed et al., '15 (submitted)]. A sliding-window of 12

¹Department of Physics and Astronomy, University of Southern California, Los Angeles, CA, USA

Full list of author information is available at the end of the article

image pairs, corresponding to 4-5 minutes, were used to generate the MBF value at each time point.

Results

Figure 1 shows MBF and heart rate (HR) response from the experiment with the lowest physiological noise, and the most stable heart rate during the initial rest period. MBF and HR increased after the onset of stress and also recovered to approximately the same values as during the initial rest period.

Conclusions

This study demonstrates the feasibility of noninvasive and continuous monitoring of MBF using myocardial ASL with mild stress. Because MBF is related to certain factors such as HR, stroke volume, oxygen demand and vessel dilation, this ASL approach could be used for continuous monitoring of both perfusion and endothelial function.





© 2015 Do et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http:// creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Funding

American Heart Association; Wallace H. Coulter Foundation.

Authors' details

¹Department of Physics and Astronomy, University of Southern California, Los Angeles, CA, USA. ²Ming Hsieh Electrical Engineering Department, University of Southern California, Los Angeles, CA, USA. ³Department of Biomedical Engineering, University of Southern California, Los Angeles, CA, USA. ⁴Department of Radiology, University of Southern California, Los Angeles, CA, USA. ⁵Department of Medicine, University of Southern California, Los Angeles, CA, USA.

Published: 3 February 2015

doi:10.1186/1532-429X-17-S1-P145

Cite this article as: Do *et al.*: **Arterial spin labeling CMR perfusion** imaging is capable of continuously monitoring myocardial blood flow during stress. *Journal of Cardiovascular Magnetic Resonance* 2015 **17**(Suppl 1):P145.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit