

**Open Access** 

## **POSTER PRESENTATION**

# Assessment of left atrial systolic dyssynchrony in paroxysmal atrial fibrillation and heart failure using cardiac magnetic resonance imaging: MESA study

Luisa A Ciuffo<sup>1,2\*</sup>, Ravi Sharma<sup>1</sup>, Mohammadali Habibi<sup>6</sup>, Bharath Ambale Venkatesh<sup>1</sup>, Boaz D Rosen<sup>1</sup>, Masamichi Imai<sup>1</sup>, Steven Shea<sup>5</sup>, Robyn McClelland<sup>4</sup>, Colin O Wu<sup>3</sup>, Susan R Heckbert<sup>4</sup>, David Bluemke<sup>3</sup>, Joao A Lima<sup>1</sup>

*From* 18th Annual SCMR Scientific Sessions Nice, France. 4-7 February 2015

### Background

Left atrial (LA) remodeling in response to cardiovascular and hemodynamic stress may precede atrial fibrillation (AF) and heart failure (HF). We hypothesized that LA systolic synchronous contraction as a functional measure of LA remodeling is deranged in patients with paroxysmal AF and HF.

#### Methods

We performed a nested case-control analysis with 1:2 matching for 39 cases of paroxysmal AF (n=28, in sinus rhythm during cardiac magnetic resonance (CMR)) and HF (n=14, AF+HF; n=3) and 78 controls with similar demographic and clinical characteristics at the baseline (Table 1). LA circumferential (short axis) and longitudinal strain rate (horizontal long axis) were measured using Multi-modality Tissue Tracking (Toshiba, Japan) from

short and long-axis cine CMR images. Circumferential LA systolic dyssynchrony among 18 LA segments (6 segments x 3 slices) was evaluated as; Standard Deviation (SD) of time to pre atrial contraction Strain rate (PreA Sr<sup>c</sup>) and Peak systolic strain rate (Peak Sra<sup>c</sup>) (Figure 1). Similarly, longitudinal LA dyssynchrony parameters (among 6 segments) were: SD-Time to pre-atrial contraction strain rate (PreA Sr<sup>L</sup>) and SD-Time to peak systolic strain rate (Peak-Sra<sup>L</sup>). Wilcoxon-rank sum test (non-parametric) or two sample t-test (parametric) were used for comparison between the groups.

#### Results

In participants during MESA exam 5 (age 74 $\pm$ 8 years, 51.4% men), systolic circumferential dyssynchrony (SD-TP-PreA Sr<sup>c</sup>, msec) was significantly higher in the cases compared to controls (45.06 vs. 28.73, p<0.010). Similarly,

Table 1 Left atrial circumferential and longitudinal systolic dyssynchrony parameters among the cases and the control group.

Parameters	Controls (n=78)			Cases (n=39) (paroxysmal AF + Heart Failure)			p-value
Longitudinal	Mean ±SD	Median	IQR	Mean ±SD	Median	IQR	
SD-TP PreA SrL, msec	36.43 ± 33.53	27.40	15.50 - 46.13	51.62 ± 33.40	39.70	32.07 - 61.78	0.001
SD-TP Peak SraL	35.92 ± 43.22	23.86	16.32 - 35.60	45.23 ± 42.04	32.50	22.98 - 44.54	0.027
Circumferential							
SD-TP PreA Src, msec	28.73±13.90	26.23	19.70 - 34.98	45.06 ± 30.25	35.49	24.84 - 57.03	0.010
SD-TP Peak Srac, msec	28.95 ± 23.1	23.04	16.46 - 31.57	36.46 ± 30.93	26.83	17.45 - 44.02	0.316
LA maximum volume/BSA	33.79 ± 9.57	31.72	28.64 - 38.86	45.02 ± 17.85	44.38	31.04 - 56.44	<0.0001
Sra (%/ms)	-1.85 ± 0.85	-1.74	-2.18: -1.45	-1.03 ± 0.61	-1.04	-1.04: -0.53	<0.0001

Participants were matched for: Age, gender, race/ethnicity.

<sup>1</sup>Johns Hopkins, Baltimore, MD, USA

Full list of author information is available at the end of the article



© 2015 Ciuffo et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http:// creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.



case group had greater longitudinal dyssynchrony than controls; SD-TP PreA  $Sr^{L}$  (51.62 vs. 36.43, p=0.001) and SD-TP-Peak  $Sra^{L}$  (45.23 vs. 35.92, p=0.027) (Table 1).

### Conclusions

Patients with paroxysmal atrial fibrillation and heart failure have significantly higher LA circumferential and longitudinal systolic dyssynchrony compared to normal controls.

#### Funding

N/A.

#### Authors' details

<sup>1</sup>Johns Hopkins, Baltimore, MD, USA. <sup>2</sup>Universidade federal da Bahia, Salvador, Brazil. <sup>3</sup>National Institutes of Health, Bethesda, MD, USA. <sup>4</sup>University of Washington, Washington, DC, USA. <sup>5</sup>Columbia University, New York, NY, USA. <sup>6</sup>Union memorial Hospital, Baltimore, MD, USA.

Published: 3 February 2015

#### doi:10.1186/1532-429X-17-S1-P322

**Cite this article as:** Ciuffo *et al.*: Assessment of left atrial systolic dyssynchrony in paroxysmal atrial fibrillation and heart failure using cardiac magnetic resonance imaging: MESA study. *Journal of Cardiovascular Magnetic Resonance* 2015 **17**(Suppl 1):P322.



BioMed Central

Submit your manuscript at www.biomedcentral.com/submit