Journal of Cardiovascular Magnetic Resonance



Meeting abstract Open Access

2091 Manganese enhanced mri demonstrates a predominant role for nNOS, not eNOS, in modulating L-Type calcium channel flux in the heart

Moriel H Vandsburger*, Brent A French, Patrick A Helm, Christopher M Kramer and Frederick H Epstein

Address: University of Virginia, Charlottesville, VA, USA

* Corresponding author

from 11th Annual SCMR Scientific Sessions Los Angeles, CA, USA. 1-3 February 2008

Published: 22 October 2008

Journal of Cardiovascular Magnetic Resonance 2008, 10(Suppl 1):A360 doi:10.1186/1532-429X-10-S1-A360

This abstract is available from: http://jcmr-online.com/content/10/S1/A360

© 2008 Vandsburger et al; licensee BioMed Central Ltd.

Introduction

Modulation of L-Type Calcium Channel (LTCC) flux plays an important role in calcium cycling and contractility. Based upon localization within the cardiomyocyte, prevailing opinion is that neuronal nitric oxide synthase (nNOS) modulates sarcoplasmic reticular calcium release, while endothelial NOS (eNOS) modulates LTCC flux. Counter to this hypothesis, a recent *in vitro* study suggests that nNOS modulates LTCC flux. Since Mn²⁺ enters the myocyte through the LTCC in proportion to Ca²⁺ flux and shortens T1, Mn-enhanced MRI may be used to probe *in vivo* LTCC flux.

Methods

Eleven wild type (WT), 8 eNOS-/- and 8 nNOS-/- mice aged 3 months were studied by Mn-enhanced MRI on a 4.7 T MRI system (Varian, CA). Two mid-ventricular short axis slices were acquired using a saturation recovery sequence with a constant repetition time of 200 ms. Images were acquired every 2 to 3 minutes for 20 minutes prior to and 45 minutes following a 30 minute infusion of MnCl₂ at a dose of 0.42 ηg/kg·min. Signal-to-noise ratio (SNR) was measured from the entire myocardium for each slice and plotted against time (Figure 1). The portion of the SNR vs. time curve corresponding to MnCl₂ infusion was isolated and the slope of the linear fit to that data was used as the LTCC index (LTCCI). Additionally, systolic blood pressure (BP) was measured using a tail cuff system.

Results

Higher BP was found in eNOS-/- mice compared to nNOS-/- (106 ± 4 WT, 111 ± 4 eNOS-/-, 94 ± 3 nNOS-/- p = 0.01 vs. eNOS-/-). Heart rate was similar between all groups (481 ± 18 WT, 470 ± 16 eNOS-/-, 490 ± 29 nNOS-/-, P = NS). LTCCI trended higher in eNOS-/- compared to WT (P = 0.1), but was nearly twice the WT rate in nNOS-/- mice (P < 0.001) (Figure 1). Additionally, LTTCI was significantly greater in nNOS-/- compared to eNOS-/- mice (P < 0.05).

Conclusion

The significantly increased rate of Mn²⁺ enhancement in nNOS^{-/-} mice represents the first *in vivo* evidence of modulation of LTCC flux by nNOS. Although eNOS^{-/-} mice showed a trend towards a higher rate of LTCC flux compared to WT, this increase is likely a response to heightened afterload. The absence of heightened BP in nNOS^{-/-} mice paired with an increased LTCCI demonstrates that nNOS, not eNOS, plays a dominant role in modulating LTCC flux.




Figure I Impact of $MnCl_2$ infusion on signal intensity within the myocardium (A). Example SNR vs. time curves for each group. Slopes of linear fits represents measures of flux rate (B). LTTCI shown to be significantly greater in $nNOS^{-J-}$ mice vs. WT (C).

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- \bullet yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing_adv.asp

