

POSTER PRESENTATION

Open Access

Serial assessment of myocardial T2 in Duchenne muscular dystrophy

Janaka Wansapura^{1*}, Kan Hor¹, Wojciech Mazur², Robert Fleck¹, Michael Taylor¹, D Woodrow Benson¹, William Gottliebson¹

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Background

Duchenne Muscular Dystrophy (DMD), a lethal X-linked skeletal and cardiac myopathy, affects 1/3500 males[1-2]. MRI studies have shown occult ventricular dysfunction and myocardial fibrosis in DMD patients. Previously we used the Full Width of Half Maximum (*FWHM*) of T2 distribution in LV to quantify the myocardial structural heterogeneity in DMD patients. In DMD subject groups, we showed that *FWHM* of the T2 histogram rose progressively with age and decreasing EF indicating that functional impairments could be associated with pre-existing abnormalities in tissue structure in young DMD patients. In this study we assessed the T2 distribution in DMD patients at two time points. We hypothesized that serial *FWHM* changes can be detected in individual DMD patients during a time when left ventricular ejection fraction (EF) changes are insignificant.

Methods

MRI Data of eighteen DMD patients obtained at two time points were analyzed (mean=12 years, range =8-18 years) with a mean time of 2.3 years between the studies. Spin echo images of the left ventricle in the short axis plane were acquired using a black blood dual spin echo method. Imaging parameters were: Slice thickness = 5mm, in plane resolution = 1.4mm×1.4mm, echo train length = 5, Echo times: TE₁ = 6 ms, TE₂ = 34ms. A histogram of LV T2 distribution with bin size equal to 1 ms was constructed for each subject. The Full Width at Half Maximum (*FWHM*) was calculated after applying box car averaging. The *FWHM* was defined as the width of the histogram at half the maximum height.

Results and conclusions

In the interval between studies the *FWHM* increased in all but two patients (89%). The average increase in *FWHM* was 6.8 ± 3.9 ms. The mean T2 increased in 5 patients while it declined in 13 patients. In a previous study we showed that *FWHM* did not change with age in normal subjects. Thus increase in T2 heterogeneity quantified by *FWHM* indicates progression of disease over a relatively short period.

Author details

¹Cincinnati Children's Hospital, Cincinnati, OH, USA. ²Christ Hospital, Cincinnati, OH, USA.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P282

Cite this article as: Wansapura et al.: Serial assessment of myocardial T2 in Duchenne muscular dystrophy. *Journal of Cardiovascular Magnetic Resonance* 2011 13(Suppl 1):P282.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Cincinnati Children's Hospital, Cincinnati, OH, USA
Full list of author information is available at the end of the article