

POSTER PRESENTATION

Open Access

Gender differences in response to cold pressor test assessed with velocity-encoded cine-MR imaging of the coronary sinus

Alexis Jacquier*, Antonin Flavian, Frank Kober, Francesca Carta, Boris Maurel, Patrick J Cozzone, Monique Bernard

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Introduction

Gender-specific differences in cardiovascular risk are well known, and current evidence supports an existing role of coronary endothelium function in these differences.

Purpose

To assess non invasively coronary endothelial function in male and female young volunteers by myocardial blood flow (MBF) measurement using coronary sinus (CS) flow quantification by velocity encoded cine MRI at rest and during cold pressor test (CPT).

Methods


Twenty-four healthy volunteers (12 men, 12 women) underwent MRI in a 3 Tesla scanner (Verio, Siemens, Erlangen, Germany). CPT was performed by immersing the right ankle in ice-water during 4 min. Heart rate and blood pressure were monitored throughout the protocol. Coronary sinus flow was measured at rest and during CPT using a non breath-hold velocity encoded phase contrast cine MRI (TR/TE: 45 ms / 2 ms, slice thickness: 5.5mm, FOV: 250², averages: 11, matrix: 256², flow encoding 70cm/s, flip angle: 25°, acquisition time: 4 min, GRAPPA k-space reduction factor: 4). Left ventricular volumes, function and morphology were evaluated using SSFP sequence. MBF was calculated combining coronary sinus flow quantification and morphologic data using Argus work station (Siemens, Erlangen, Germany). Coronary endothelial function was assessed by comparing MBF at rest and during CPT. Coronary vascular resistance (CVR), rate pressure product and endothelium-dependent vasodilatation index (EDVI) were calculated.

Results

At baseline, MBF was $0.63 \pm 0.23 \text{ mL}\cdot\text{g}^{-1}\cdot\text{min}^{-1}$ in men and $0.79 \pm 0.21 \text{ mL}\cdot\text{g}^{-1}\cdot\text{min}^{-1}$ in women ($P=\text{ns}$) (figure 1). During CPT, the rate pressure product in men significantly increased by $49 \pm 36 \%$ ($p<0.0001$) and in women by $52 \pm 22 \%$ ($p<0.0001$). MBF increased significantly in both men and women by $0.22 \pm 0.19 \text{ mL}\cdot\text{g}^{-1}\cdot\text{min}^{-1}$ ($p=0.0022$) and by $0.73 \pm 0.43 \text{ mL}\cdot\text{g}^{-1}\cdot\text{min}^{-1}$ ($p=0.0001$), respectively. The increase in MBF was significantly higher in women than in men ($p = 0.0012$).

Conclusions

MRI coronary sinus flow quantification for measuring myocardial blood flow revealed a higher response of MBF to CPT in women than in men. This finding may reflect gender differences in endothelial-dependent



vasodilatation. This non invasive rest/stress protocol may become helpful to study endothelial function in normal physiology and in physiopathology.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P346

Cite this article as: Jacquier *et al.*: Gender differences in response to cold pressor test assessed with velocity-encoded cine-MR imaging of the coronary sinus. *Journal of Cardiovascular Magnetic Resonance* 2011 13(Suppl 1):P346.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

