

POSTER PRESENTATION

Open Access

Relationship between proximal aortic stiffness assessed with CMR and left ventricular diastolic function in a bicentric asymptomatic population with preserved left ventricular ejection fraction

Alban Redheuil^{1*}, Elie Mousseaux¹, Wen-Chung Yu², Nadja Kachenoura¹, Ludivine Perdrix¹, Elzbieta Chamera³, David Bluemke⁴, Joao Lima³

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Objective

To evaluate the direct relation between proximal aortic function and left ventricular diastolic function in individuals with normal LVEF free of overt cardiovascular disease.

Background

Left ventricular (LV) diastolic dysfunction is considered to be an important determinant of heart failure with preserved LV ejection fraction (LVEF). Hypertension, diabetes and aging leading to increased aortic stiffness have been related to impaired diastolic function. Novel CMR measures of proximal aortic function have been recently shown to be better markers of vascular aging than conventional measures of arterial stiffness but their relationship to LV diastolic function has not been reported.

Methods

We studied 144 subjects (70 men, 74 women, age 44 ± 16 [13-79] years) from 2 academic centers (Baltimore, USA; Paris, France). Ascending aortic strain was determined by CMR using an automated segmentation of SSFP cine acquisitions. Aortic arch pulse wave velocity (PWV) was measured by CMR from phase-contrast sequence for flow-wave analysis and multiple T1 spin echo images for aortic length measurement. Conventional LV diastolic function indices were measured by echo-Doppler (E/A, e' , E/e'); carotid pressures and carotid-femoral PWV

(cfPWV) were measured using applanation tonometry. Central pressures were used to calculate aortic distensibility and carotid augmentation index (AIx).

Results

As shown on Figure 1, increased aortic arch stiffness was associated with decreased E/A ratio, decreased mitral annular e' velocity and increased E/e' ratio. In multivariate linear regression analysis we found an independent association between impaired ascending aortic distensibility and decreased E/A ratio ($p=0.01$, $R^2=0.42$) and e' velocities ($p=0.006$, $R^2=0.62$). In addition, we found an independent association between increased aortic arch PWV and increased E/e' ratio ($p<0.001$, $R^2=0.46$). These relationships were independent of age, gender, body mass index and mean central blood




Figure 1 Relationship between E/A and E/e' ratios and aortic arch PWV

¹European Hospital Georges Pompidou and INSERMU678, Paris, France
Full list of author information is available at the end of the article

pressure. Moreover, these associations remained significant after further adjustment for LV mass. Similar trends were observed using central pulse pressure, AIx and cfPWV as markers of arterial stiffness but reached lower degrees of statistical significance compared to proximal markers of aortic function.

Conclusions

Proximal aortic function measured by CMR is strongly associated with subclinical impairment in diastolic left ventricular function in subjects with preserved LVEF independent of the effects of age, gender, BMI, mean central pressure and LV mass. These novel markers of proximal aortic stiffness estimated by CMR have a stronger correlation with LV diastolic function than conventional markers of arterial stiffness.

Author details

¹European Hospital Georges Pompidou and INSERMU678, Paris, France.

²Taipei Veterans General Hospital, Division of Cardiology, Taipei, Taiwan.

³Johns Hopkins University, Baltimore, MD, USA. ⁴Radiology and Imaging Sciences, National Institutes of Health, Bethesda, MD, USA.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P378

Cite this article as: Redheuil *et al.*: Relationship between proximal aortic stiffness assessed with CMR and left ventricular diastolic function in a bicentric asymptomatic population with preserved left ventricular ejection fraction. *Journal of Cardiovascular Magnetic Resonance* 2011 13(Suppl 1):P378.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

