

ORAL PRESENTATION

Open Access

Longitudinal monitoring of cardiac dysfunction with MRI in a mouse model of obesity and type 2 diabetes

Inès Abdesselam^{1,2*}, Thomas Troalen¹, Pauline Pepino¹, Anne Dutour^{2,3}, Frank Kober¹, Monique Bernard¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Obesity consequences include type 2 diabetes and cardiovascular diseases. Ectopic fat deposition within certain organs such as heart and liver, have recently been shown to be important independent risk factors for the development of these diseases. The purpose of this study was to perform a longitudinal analysis of the changes in cardio-metabolic functions associated with adiposity increase and insulin resistance development.

Methods

Male C57BL6R mice were subjected to a high fat high sucrose diet (HFHSD, N = 10) or a regular show diet (RD, N = 10) during 4 months in order to induce obesity and insulin resistance. MRI was performed on a Bruker Biospec AVANCE 4.7T system for animal examination. Mice underwent complete exploration every month as follows: 1/determination of body fat content and distribution including ectopic fat using MRS with PRESS sequence and body imaging, 2/cardiac function assessment with 4-cavity and short-axis cine sequences, and 3/cardiac perfusion investigation with arterial spin labeling [1].

Results

Mice fed with HFHSD became significantly obese compared to mice fed with RD (47.3 ± 0.7 g vs. 32.9 ± 0.1 g). After 4 months of HFHSD, obese mice developed cardiac dysfunctions. Indeed, they presented a significant decrease in end diastolic volume (EDV) compared to control mice (66.9 ± 3.7 mL vs. 85.8 ± 3.0 mL, $p = 0.001$), stroke volume

(SV) (47.4 ± 2.8 mL vs. 58.5 ± 2.8 mL $p = 0.01$). A significant decrease in cardiac perfusion (7.0 ± 0.7 mL/g/min vs. 9.1 ± 0.6 mL/g/min, $p = 0.03$) was also observed (Figure 1). These anomalies have been developed subsequently to a significant triglycerides accumulation in heart ($1.1 \pm 0.20\%$ vs. $0.7 \pm 0.08\%$, $p = 0.02$) and liver ($5.0 \pm 1.3x\%$ vs. $1.8 \pm 0.09\%$, $p = 0.002$) which appears earlier in the 1st month (Figure 2). Furthermore, cardiac perfusion was significantly associated to myocardial ($r = -0.5$, $p = 0.04$) and hepatic ($r = -0.64$, $p = 0.008$) triglyceride content at 4 months although these two ectopic fats are not correlated together. Interestingly, cardiac perfusion at 4 months was also associated to hepatic triglyceride content at 3 months ($r = -0.56$, $p = 0.03$), while no change in cardiac perfusion was still observed. Moreover, increased hepatic triglyceride content was correlated to myocardial function decreases at 4 months: EDV ($r = -0.65$, $p = 0.008$), SV ($r = -0.64$, $p = 0.01$).

Conclusions

These preliminary data established the chronological order of anomalies appearance associated to diet-induced obesity and insulin resistance in a 4 month longitudinal assessment. Hepatic triglyceride content seems to be a relevant predictive factor of cardiac anomalies. This experimental protocol might be useful to assess the effect of anti-diabetic drugs on cardiac function.

Funding

QASAREM.

¹Aix-Marseille University, CRMBM/CEMEREM CNRS 7339, Marseille, France
Full list of author information is available at the end of the article




Figure 1




Figure 2

Authors' details

¹Aix-Marseille University, CRMBM/CEMEREM CNRS 7339, Marseille, France.

²Aix-Marseille University, NORT, Inserm 1062, Inra 1260, Marseille, France.

³Endocrinology, Metabolic Disease and Nutrition, CHU Nord, Marseille, France.

Published: 16 January 2014

Reference

1. Cine-ASL: A steady-pulsed arterial spin labeling method for myocardial perfusion mapping in mice. Part I. Experimental study. In *Magn Reson Med* Troalen T, Capron T, Cozzone PJ, Bernard M, Kober F 2013.

doi:10.1186/1532-429X-16-S1-O113

Cite this article as: Abdesselam *et al.*: Longitudinal monitoring of cardiac dysfunction with MRI in a mouse model of obesity and type 2 diabetes. *Journal of Cardiovascular Magnetic Resonance* 2014 **16** (Suppl 1):O113.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

