

POSTER PRESENTATION

Open Access

Left ventricular dyssynchrony can be observed via cine CMR with use of aortic valve timing

Francisco Contijoch*, Kelly Rogers, Walter R Witschey, Robert C Gorman, Yuchi Han

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Cine CMR is the gold standard for evaluation of left ventricular function with high spatial and temporal resolution of the EKG-gated acquisition. Measurements of dyssynchrony have previously been performed using myocardial tagging to measure relative delays in peak shortening of opposing walls. We aim to utilize non-tagged images to quantify dyssynchrony in clinical patients. This is performed by combining aortic valve timing information from a left ventricular outflow tract (LVOT) image series with segmentations of short axis datasets.

Methods

Short-axis stacks and LVOT images were acquired using standard retrospective EKG-gated bSSFP cine imaging in 63 clinical patients with a range of clinical conditions (mean EF = 44.5 ± 19.0%) as well as 29 patients with prolonged QRS duration. Scan parameters were as follows: in-plane resolution: 1.25 - 2.08 mm, slice spacing: 8 mm with 2 mm gap, reconstructed temporal resolution: 18.2 - 58.8 ms. Aortic valve opening and closing as a percentage of cardiac cycle was identified on LVOT images. Volumetric evaluation of short axis images was performed via semi-automated segmentation for all phases and all left ventricular slices (ITK-SNAP, Philadelphia PA). Dyssynchrony was measured by the standard deviation of the timing difference between each slice minimum area to aortic valve closure as expressed by the equation in Figure 1 where $\tau_{\text{slice min}}$ corresponds to the phase at which a slice achieves its minimum area.

Results

The use of patients with prolonged QRS complexes allows for separation of measured dyssynchrony based on

both EF and QRS. Patients were categorized into low EF ($\leq 35\%$), mid EF ($35\% < \text{EF} < 50\%$) and normal EF ($\geq 50\%$). The results are shown in Figure 1. As has been observed using tagged-MRI, patients with low EF demonstrate higher dyssynchrony than both EF-matched controls as well as patients with long QRS but normal EF.

Conclusions

This technique allows for a slice-by-slice timing based regional evaluation of dyssynchrony. In addition, subsequent analysis of particular dyssynchronous slices could be performed to provide additional regional information.

$$\text{Dyssynchrony} = \sqrt{\frac{1}{\text{slices}} \sum_{\text{slices}} (\tau_{\text{slice min}} - \tau_{\text{aortic closure}})^2}$$




Figure 1 End-systolic dyssynchrony was quantified in 67 patients with a range of EF and QRS durations. Observed values were classified based on high, mid and low ED as well as normal prolonged QRS duration. The observed dyssynchrony for patients with low EF and prolonged QRS is higher than all other groups. However, more patients are necessary to achieve significance.

Funding

K99-HL108157, R01-HL103723, T32HL007954, T32-EB009384.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P243

Cite this article as: Contijoch *et al.*: Left ventricular dyssynchrony can be observed via cine CMR with use of aortic valve timing. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P243.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

