



POSTER PRESENTATION

Open Access

Cardiovascular magnetic resonance is feasible in many patients aged 3 to 8 years without general anesthesia or sedation

Ahmed E Kharabish^{1*}, Naira Mkrtchyan¹, Christian Meierhofer¹, Stefan Martinoff², Peter Ewert¹, Heiko Stern¹, Sohrab Fratz¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Cardiovascular Magnetic Resonance (CMR) of pediatric patients has become routine clinical practice. Patients under eight years are usually examined under general anesthesia (GA) or sedation without intubation. However, in our clinical experience, CMR may be feasible in patients aged 3 to 8 years without general anesthesia or sedation without intubation.

Methods

Retrospectively studied datasets of total number of 71 patients aged between 3 and 8 years. The total cohort was divided into two groups, first with no general anesthesia or sedation without intubation (no GA or sedation) and the second group patients with general anesthesia or sedation without intubation (GA or sedation). The patients' age groups and scan durations for each group, percentage of successfully answering the clinical question in each group, total number of scanned sequences in each group, and number of sequences per study were recorded and compared between both groups.

Results

Forty-four patients in the no GA or sedation group, 27 in the GA or sedation group. The scan duration was in the no GA or sedation group: 35 minutes \pm 20.44 minutes, and in the GA or sedation group: 60 minutes \pm 31 minutes ($p < 0.001$). The percentage of successful reports was 95% (42 of 44) in the no GA or sedation group and 89% (24 of 27) in the patients with GA or sedation group.

Conclusions

In scope of understanding the CHD hemodynamics, and clinical requests and questions; the decision of using neither general anesthesia with intubation nor sedation with without intubation was favored by our center's CMR unit over examining the patients under general anesthesia or sedation, to avoid the effect of general anesthesia and sedation on cardiac function during CMR, decrease the CMR scan duration, and increase the feasibility of the CMR in the young age group between three and eight years.

Funding

All authors have no conflict of interest.

Authors' details

¹Pediatric Cardiology and Congenital Heart Defects, German Heart Center, Munich, Germany. ²Radiology and Nuclear Medicine, German Heart Center, Munich, Germany.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P377

Cite this article as: Kharabish et al.: Cardiovascular magnetic resonance is feasible in many patients aged 3 to 8 years without general anesthesia or sedation. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P377.

¹Pediatric Cardiology and Congenital Heart Defects, German Heart Center, Munich, Germany

Full list of author information is available at the end of the article