

POSTER PRESENTATION

Open Access

# Assessing right ventricular function in patients with pulmonary hypertension based on noninvasive measurements: correlation between cardiac MRI, ultrasonic cardiogram, multidetector CT and right heart catheterization

Xiaojuan Guo<sup>1\*</sup>, Min Liu<sup>1</sup>, Zhanhong Ma<sup>1</sup>, tao jiang<sup>1</sup>, Yuanhua Yang<sup>2</sup>, zhenguo zhai<sup>2</sup>, Renyou zhai<sup>1</sup>, Tianjing Zhang<sup>3</sup>

From 18th Annual SCMR Scientific Sessions  
Nice, France. 4-7 February 2015

## Background

The assessment of right ventricular function is of great importance in the management of patients with pulmonary hypertension (PH). Our aim is to compare the value of cardiac MRI (CMRI), ultrasonic cardiogram (CUS), multidetector CT (MDCT) in assessing right ventricular function, and to evaluate the correlation between parameters derived by MRI, CUS, MDCT and the indicators of right ventricular function derived by right heart catheterization (RHC) in patients with PH.

## Methods

Thirty one consecutive patients with PH (17 males and 14 females; mean age,  $54.61 \pm 11.74$  years; range, 33 to 76 years) were prospectively enrolled. All patients underwent RCH to get hemodynamic parameters. Then, calculations including pulmonary vascular resistance index (PVRI), right cardiac work index (RCWI), right ventricular stroke work index (RVSWI) were performed. All patients underwent CMRI to get parameters including right ventricular end-systolic volume (ESV), end-diastolic dimension (EDV), stroke volume (SV), ejection fraction (EF), the cardiac muscle mass (CMS). 28 patients underwent CUS to get these parameters including right ventricular Tei index, right ventricular fractional area change (RV FAC), RV-ESV, RV-EDV. And 25 patients

underwent MDCT to get these parameters including right /left ventricular internal diameter (RVd/LVd), right /left ventricular diastole maximum area (RVa/LVa), cobb angle, right ventricular free wall thickness (RVWT). All examinations were executed within 7 days. These parameters obtained by MRI, CUS and MDCT were correlated with those of RCH respectively by Spearman or Pearson correlation analysis.

## Results

Most parameters of RV function derived by CMRI, CUS, MDCT correlated moderately with mPAP, PVRI. mPAP had strong correlation with MRI-ventricular mass index (VMI) ( $r=0.528$ ,  $P=0.002$ ), RVWT ( $r=0.554$ ,  $P=0.005$ ); PVRI had strong correlation with MRI-EF ( $r=-0.647$ ,  $P=0.000$ ), MRI-VMI ( $r=-0.567$ ,  $P=0.001$ ), CT-RVd ( $r=0.536$ ,  $P=0.006$ ), RVd/LVd ( $r=0.530$ ,  $P=0.006$ ), CT-Cobb angle ( $r=0.608$ ,  $P=0.001$ ), RVWT ( $r=0.586$ ,  $P=0.003$ ). Table 1 showed the correlation of right ventricular function between CMRI, CUS, MDCT and RHC. RHC-SV had the strong correlation with MRI-EF ( $r=0.557$ ,  $P=0.001$ ), CT-RVd ( $r=-0.502$ ,  $P=0.011$ ), CT-Cobb angle ( $r=-0.503$ ,  $P=0.003$ ).

## Conclusions

The parameters obtained by MRI, CUS and MDCT had moderate correlation with RHC-SV. MRI-EF was the best parameter to reflect RV function which could be used to exactly evaluate RV function in patient with PH.

<sup>1</sup>Department of Radiology, Beijing Chao-Yang Hospital, Capital Medical University, Beijing, China

Full list of author information is available at the end of the article

**Table 1 Correlation coefficients of right ventricular function parameters between CMRI, CUS, M DCT and RHC.**

	6MWD		RHC-SV		RCWI		RVSWI	
	r	P	r	P	r	P	r	P
MRI-SV(n=31)	0.073	0.713	0.457	0.010	0.322	0.077	0.324	0.075
MRI-EF	0.473	0.011	0.557	0.001	0.389	0.031	0.307	0.093
MRI-VMI	-0.472	0.011	-0.444	0.012	-0.259	0.159	-0.173	0.351
US-SV(n=28)	-0.268	0.186	0.016	0.937	-0.201	0.305	-0.071	0.718
US-EF	-0.019	0.923	0.406	0.026	0.015	0.939	0.023	0.905
US-Tei	-0.175	0.448	-0.480	0.024	-0.387	0.075	-0.419	0.042
US-RVFAC	0.046	0.822	0.385	0.043	0.289	0.136	0.199	0.311
CT-RVd(n=25)	-0.373	0.079	-0.502	0.011	-0.486	0.014	-0.468	0.018
CT-RVa	-0.351	0.109	-0.488	0.016	-0.342	0.102	-0.362	0.082
CT-Cobb angle	-0.569	0.004	-0.503	0.003	-0.430	0.025	-0.395	0.041
CT-RVWT	-0.098	0.666	-0.279	0.186	-0.024	0.911	0.023	0.914

6MWD, 6-min walking distance; EF, ejection fraction; VMI, ventricular mass index; RCWI, right cardiac work index; RHC, right heart catheterization; RVa, right ventricular diastole maximum area; RVd, right ventricular internal diameter; RVFAC, right ventricular fractional area change; RVSWI, right ventricular stroke work index; RVWT, right ventricular free wall thickness; SV, stroke volume; Tei, Tie index; P<0.05.

Among these noninvasive measurements, CMRI is the optimal method to assess RV function, and then is the MDCT and the last is CUS.

## Funding

the National 12th Five-Year Researching Foundation (2011BAI11B17).

## Authors' details

<sup>1</sup>Department of Radiology, Beijing Chao-Yang Hospital, Capital Medical University, Beijing, China. <sup>2</sup>Respiratory Diseases Researching Center of Capital Medical University, Beijing, China. <sup>3</sup>Siemens Ltd., China, Beijing, China.

Published: 3 February 2015

doi:10.1186/1532-429X-17-S1-P185

**Cite this article as:** Guo et al.: Assessing right ventricular function in patients with pulmonary hypertension based on noninvasive measurements: correlation between cardiac MRI, ultrasonic cardiogram, multidetector CT and right heart catheterization. *Journal of Cardiovascular Magnetic Resonance* 2015 17(Suppl 1):P185.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

