

CORRECTION

Open Access



# Correction to: Subclinical myocardial injury in patients with Facioscapulohumeral muscular dystrophy 1 and preserved ejection fraction – assessment by cardiovascular magnetic resonance

Edyta Blaszczyk<sup>1,2</sup>, Ulrike Grieben<sup>3</sup>, Florian von Knobelsdorff-Brenkenhoff<sup>1,2,4</sup>, Peter Kellman<sup>5</sup>, Luisa Schmacht<sup>1,2</sup>, Stephanie Funk<sup>1,2</sup>, Simone Spuler<sup>3</sup> and Jeanette Schulz-Menger<sup>1,2\*</sup>

## Correction to: J Cardiovasc Magn

<https://doi.org/10.1186/s12968-019-0537-4>

In the original version of this article [1], published on 29 April 2019, there is 1 error in the ‘Method’ section of the article.

The original publication states:

- A sampling protocol with reduced sensitivity to heart rate was applied [15] before and 15 min after contrast media (0.15 mmol/kg body weight **Gadoteriol**) application (T1 native: 5 s(3 s)3 s; T1 post-contrast: 4 s(1 s)3 s(1 s)2 s; TE 1.1 ms and slice thickness 6 mm).

However, the contrast media was **Gadobutrol**

The corrected information is also repeated below:

- A sampling protocol with reduced sensitivity to heart rate was applied [15] before and 15 min after contrast media (0.15 mmol/kg body weight **Gadobutrol**) application (T1 native: 5 s(3 s)3 s; T1 post-contrast: 4 s(1 s)3 s(1 s)2 s; TE 1.1 ms and slice thickness 6 mm).

## Author details

<sup>1</sup>Working Group on Cardiovascular Magnetic Resonance, Experimental and Clinical Research Center a joint cooperation between the Charité – Universitätsmedizin Berlin, Department of Internal Medicine and Cardiology and the Max-Delbrueck Center for Molecular Medicine, and HELIOS Klinikum Berlin Buch, Department of Cardiology and Nephrology, Berlin, Germany. <sup>2</sup>DZHK (German Centre for Cardiovascular Research), partner site Berlin, Berlin, Germany. <sup>3</sup>Muscle Research Unit, Experimental and Clinical Research Center a joint cooperation between the Charité Medical Faculty and the Max-Delbrueck Center for Molecular Medicine, Berlin, Germany. <sup>4</sup>Department of Cardiology, Clinic Agatharied, University of Munich, Hausham, Germany. <sup>5</sup>National Heart, Lung and Blood Institute, National Institute of Health, Berlin, Germany.

Received: 29 April 2019 Accepted: 17 May 2019

Published online: 03 June 2019

## Reference

1. Blaszczyk E, Grieben U, von Knobelsdorff-Brenkenhoff F, et al. Subclinical myocardial injury in patients with Facioscapulohumeral muscular dystrophy 1 and preserved ejection fraction – assessment by cardiovascular magnetic resonance. *J Cardiovasc Magn Reson*. 2019;21:25 <https://doi.org/10.1186/s12968-019-0537-4>.

\* Correspondence: [jeanette.schulz-menger@charite.de](mailto:jeanette.schulz-menger@charite.de)

<sup>1</sup>Working Group on Cardiovascular Magnetic Resonance, Experimental and Clinical Research Center a joint cooperation between the Charité – Universitätsmedizin Berlin, Department of Internal Medicine and Cardiology and the Max-Delbrueck Center for Molecular Medicine, and HELIOS Klinikum Berlin Buch, Department of Cardiology and Nephrology, Berlin, Germany  
<sup>2</sup>DZHK (German Centre for Cardiovascular Research), partner site Berlin, Berlin, Germany

Full list of author information is available at the end of the article

